ELSEVIER

Gene 176 (1996) 275–277

GENE
AN INTERNATIONAL JOURNAL ON GENES AND GENOMES

Author Index

Aínsa, J.A., Martín, C., Cabeza, M., De la Cruz, F. and Mendiola, M.V.
Construction of a family of *Mycobacterium/Escherichia coli* shuttle vectors derived from pAL5000 and pACYC184: their use for cloning an antibiotic-resistance gene from *Mycobacterium fortuitum* (176) 23
Alves, G., Tatro, A. and Fanning, T.
Differential methylation of human *LINE-1* retrotransposons in malignant cells (176) 39
An, Z., Farman, M.L., Budde, A., Taura, S. and Leong, S.A.
New cosmid vectors for library construction, chromosome walking and restriction mapping in filamentous fungi (176) 93
Andersen, A.B., see Braibant, M. (176) 171
Annis, S.L., see Panaccione, D.G. (176) 103
Anrather, J., see Winkler, H. (176) 67

Bach, F.H., see Winkler, H. (176) 67
Baeuerle, P.A., see Linker, R.A. (176) 119
Battista, M.C., Bergamini, G., Campanini, F., Landini, M.P. and Ripalti, A.
Intracellular production of a major cytomegalovirus antigenic protein in the methylotrophic yeast *Pichia pastoris* (176) 197
Baulard, A., Kremer, L., Supply, P., Vidaud, D., Bidart, J.-M., Bellet, D. and Locht, C.
A new series of mycobacterial expression vectors for the development of live recombinant vaccines (176) 149
Bavoil, P.M., see Hsia, R.-c. (176) 155
Bavoil, P.M., see Hsia, R.-c. (176) 163
Bellet, D., see Baulard, A. (176) 149
Bellitti, M., see Rocco, L. (176) 185
Bergamini, G., see Battista, M.C. (176) 197
Bhat, S.P., see Ruiz, A. (176) 237
Bidart, J.-M., see Baulard, A. (176) 149
Biegel Carson, S.D., Thomas, C.E. and Elkins, C.
Cloning and sequencing of a *Haemophilus ducreyi fur* homolog (176) 125
Bird, P., see Niforas, P. (176) 139
Blanco, J., José, J., Coque, R. and Martín, J.F.
Characterization of the *secA* gene of *Streptomyces lividans* encoding a protein translocase which complements an *Escherichia coli* mutant defective in the ATPase activity of SecA (176) 61
Bocchini, V., see De Vendittis, E. (176) 27
Bok, D., see Ruiz, A. (176) 237
Bolívar, J., Goenechea, L.G., Grenett, H., Pendón, C. and Valdivia, M.M.
Cloning and sequencing of the genes encoding the hamster ribosomal transcription factors UBF1 and UBF2[1] (176) 257
Bordonné, R. and Tarassov, I.
The yeast *SME1* gene encodes the homologue of the human E core protein (176) 111
Bose, S.K., see Gao, B. (176) 269
Bowman, T.L., Kaludov, N.K., Klein, M. and Hurt, M.M.
An *H3* coding region regulatory element is common to all four nucleosomal classes of mouse histone-encoding genes (176) 1
Braibant, M., Lefèvre, P., De Wit, L., Peirs, P., Ooms, J., Huygen, K., Andersen, A.B. and Content, J.
A *Mycobacterium tuberculosis* gene cluster encoding proteins of a phosphate transporter homologous to the *Escherichia coli* Pst system (176) 171
Brostjan, C., see Winkler, H. (176) 67
Budde, A., see An, Z. (176) 93

Cabeza, M., see Aínsa, J.A. (176) 23
Campanini, F., see Battista, M.C. (176) 197
Carbonell, X. and Villaverde, A.
Peptide display on functional tailspike protein of bacteriophage P22 (176) 225
Chauthaiwale, V.M., see Gao, C.Y. (176) 243
Chen, W., see Pasternak, C. (176) 177
Chen, Y.-P., see Lee, Y.-J. (176) 85
Chenal, V., Souque, P., Markovits, A. and Griffais, R.
Choosing highly specific primers for the polymerase chain reaction using the octomer frequency disparity method: application to *Chlamydia trachomatis* (176) 97
Chow, W.-Y., see Lee, Y.-J. (176) 85
Chu, M.D., see Niforas, P. (176) 139
Clarke, A.K., see Hiltonen, T. (176) 263
Content, J., see Braibant, M. (176) 171
Coque, R., see Blanco, J. (176) 61
Csizmadia, V., see Winkler, H. (176) 67

De la Cruz, F., see Aínsa, J.A. (176) 23
De Vendittis, E. and Bocchini, V.
Protein-encoding genes in the sulfothermophilic archaea *Sulfolobus* and *Pyrococcus* (176) 27
De Wit, L., see Braibant, M. (176) 171
Dixon, N.E., see Love, C.A. (176) 49
Dorner, F., see Pfleiderer, M. (176) 131

Elbrecht, A., see Neeper, M. (176) 203
Elkins, C., see Biegel Carson, S.D. (176) 125

Falkner, F.G., see Pfleiderer, M. (176) 131
Fanning, T., see Alves, G. (176) 39
Farman, M.L., see An, Z. (176) 93
Flores, S.C., see Gao, B. (176) 269
Fukuchi, T., see Tsuchiya, E. (176) 35

Gagnon, F., see Islam, N. (176) 9
Galuska, S., see Neeper, M. (176) 203
Gao, B., Flores, S.C., Bose, S.K. and McCord, J.M.
A novel *Escherichia coli* vector for oxygen-inducible high level expression of foreign genes (176) 269
Gao, C.Y., Chauthaiwale, V.M., Rampalli, A.M. and Zelenka, P.S.
Expression of alternatively spliced PCTAIRE-1 mRNA in PC12 cells and neonatal rat brain (176) 243
Ghosh, S. and Lowenstein, J.M.
A multifunctional vector system for heterologous expression of proteins in *Escherichia coli* (176) 249
Gilbert, M., see Ostiguy, S. (176) 265
Gobright, E., see Mertens, B. (176) 273
Goenechea, L.G., see Bolívar, J. (176) 257
Grenett, H., see Bolívar, J. (176) 257
Griffais, R., see Chenal, V. (176) 97

Hamada, T., Tashiro, K., Tada, H., Inazawa, J., Shirozu, M., Shibahara, K., Nakamura, T., Martina, N., Nakano, T. and Honjo, T.
Isolation and characterization of a novel secretory protein, stromal cell-derived factor-2 (SDF-2) using the signal sequence trap method (176) 211
Heck, C., see Pasternak, C. (176) 177
Hiltonen, T., Clarke, A.K., Karlsson, J. and Samuelsson, G.
A cDNA coding for glutathione *S*-transferase from the unicellular green algae *Coccomyxa* sp.[1] (176) 263
Hofmann, K.J., see Neeper, M. (176) 203
Honjo, T., see Hamada, T. (176) 211
Hove-Jensen, B., see Krath, B.N. (176) 73
Hsia, R.-c. and Bavoil, P.M.
Sequence analysis of the *omp2* region of *Chlamydia psittaci* strain GPIC: structural and functional implications (176) 155
Hsia, R.-c. and Bavoil, P.M.
Homologs of *Escherichia coli recJ*, *gltX* and of a putative 'early' gene of avian *Chlamydia psittaci* are located upstream of the 'late' *omp2* locus of *Chlamydia psittaci* strain guinea pig inclusion conjunctivitis (176) 163
Hurt, M.M., see Bowman, T.L. (176) 1
Huygen, K., see Braibant, M. (176) 171

Inazawa, J., see Hamada, T. (176) 211
Islam, N., Poitras, L., Gagnon, F. and Moss, T.
Antisense and sense poly(A)$^-$RNAs from the *Xenopus laevis* pyruvate dehydrogenase gene loci are regulated with message production during embryogenesis (176) 9

José, J., see Blanco, J. (176) 61

Kakikawa, M., see Oki, M. (176) 215
Kaltschmidt, C., see Linker, R.A. (176) 119
Kaludov, N.K., see Bowman, T.L. (176) 1
Kambara, H., see Okano, K. (176) 231
Karlsson, J., see Hiltonen, T. (176) 263
Klein, M., see Bowman, T.L. (176) 1
Kluepfel, D., see Ostiguy, S. (176) 265
Klug, G., see Pasternak, C. (176) 177
Kodaira, K.-I., see Oki, M. (176) 215
Kodama, K., see Shimizu, E. (176) 45
Konopiński, R., Nowak, R. and Siedlecki, J.A.
Alternative polyadenylation of the gene transcripts encoding a rat DNA polymerase β (176) 191
Krath, B.N. and Hove-Jensen, B.
Bacillus caldolyticus prs gene encoding phosphoribosyl-diphosphate synthase (176) 73
Kremer, L., see Baulard, A. (176) 149
Kuo, J.S., Veale, R., Maxwell, B. and Sive, H.
Translational inhibition by 5′-polycytidine tracts in *Xenopus* embryos and in vitro (176) 17
Kurano, K., see Tsuchiya, E. (176) 35

Lamont, I.L., see McMorran, B.J. (176) 55
Landini, M.P., see Battista, M.C. (176) 197
Lee, Y.-J., Chen, Y.-P., Wang, S.-H., Chow, W.-Y. and Lin, L.-Y.
Structure and expression of metallothionein gene in ducks (176) 85
Lefèvre, P., see Braibant, M. (176) 171
Leong, S.A., see An, Z. (176) 93
Lilley, P.E., see Love, C.A. (176) 49
Lin, L.-Y., see Lee, Y.-J. (176) 85
Linker, R.A., Baeuerle, P.A. and Kaltschmidt, C.
Cloning of the murine relA (p65 NF-κB) gene and comparison to the human gene reveals a distinct first intron (176) 119
Locht, C., see Baulard, A. (176) 149
Love, C.A., Lilley, P.E. and Dixon, N.E.
Stable high-copy-number bacteriophage λ promoter vectors for overproduction of proteins in *Escherichia coli* (176) 49
Lowe, R., see Neeper, M. (176) 203
Lowenstein, J.M., see Ghosh, S. (176) 249

Markovits, A., see Chenal, V. (176) 97
Martín, C., see Aínsa, J.A. (176) 23
Martín, J.F., see Blanco, J. (176) 61
Martina, N., see Hamada, T. (176) 211
Matsuzaki, G., see Tsuchiya, E. (176) 35
Maxwell, B., see Kuo, J.S. (176) 17
McCord, J.M., see Gao, B. (176) 269
McMorran, B.J., Merriman, M.E., Rombel, I.T. and Lamont, I.L.
Characterisation of the *pvdE* gene which is required for pyoverdine synthesis in *Pseudomonas aeruginosa* (176) 55
Mendiola, M.V., see Aínsa, J.A. (176) 23
Merriman, M.E., see McMorran, B.J. (176) 55
Mertens, B., Gobright, E. and Seow, H.-F.
The nucleotide sequence of the bovine interleukin-5-encoding cDNA[1] (176) 273
Miyakawa, T., see Tsuchiya, E. (176) 35
Montano, X., see Shamsher, M. (176) 259
Moriya, S. and Ogasawara, N.
Mapping of the replication origin of the *Bacillus subtilis* chromosome by the two-dimensional gel method (176) 81
Morosoli, R., see Ostiguy, S. (176) 265
Moss, T., see Islam, N. (176) 9

Nakamura, T., see Hamada, T. (176) 211
Nakano, T., see Hamada, T. (176) 211
Natarajan, G., see Winkler, H. (176) 67
Neeper, M., Lowe, R., Galuska, S., Hofmann, K.J., Smith, R.G. and Elbrecht, A.
Molecular cloning of an avian anti-Müllerian hormone homologue (176) 203
Niforas, P., Chu, M.D. and Bird, P.
A retinoic acid/cAMP-responsive enhancer containing a cAMP responsive element is required for the activation of the mouse thrombomodulin-encoding gene in differentiating F9 cells (176) 139
Nowak, R., see Konopiński, R. (176) 191

Ogasawara, N., see Moriya, S. (176) 81
Okano, K. and Kambara, H.
Fragment walking for long DNA sequencing by using a library as small as 16 primers (176) 231
Oki, M., Kakikawa, M., Yamada, K., Taketo, A. and Kodaira, K.-I.
Cloning, sequence analysis, and expression of the genes encoding lytic functions of Bacteriophage ϕg1e (176) 215
Ooms, J., see Braibant, M. (176) 171
Ostiguy, S., Gilbert, M., Shareck, F., Kluepfel, D. and Morosoli, R.
Cloning and sequencing of the *secY* homolog from *Streptomyces lividans* 1326 (176) 265

Panaccione, D.G., Pitkin, J.W., Walton, J.D. and Annis, S.L.
Transposon-like sequences at the *TOX2* locus of the plant-pathogenic fungus *Cochliobolus carbonum* (176) 103
Pasternak, C., Chen, W., Heck, C. and Klug, G.
Cloning, nucleotide sequence and characterization of the *rpoD* gene encoding the primary sigma factor of *Rhodobacter capsulatus* (176) 177
Peirs, P., see Braibant, M. (176) 171
Pendón, C., see Bolívar, J. (176) 257
Pfleiderer, M., Falkner, F.G. and Dorner, F.
Deletion of the myristylation signal allows high-level production of the hepatitis B virus large surface glycoprotein preS1 with vaccinia virus recombinants (176) 131

Pitkin, J.W., see Panaccione, D.G. (176) 103
Poitras, L., see Islam, N. (176) 9

Rampalli, A.M., see Gao, C.Y. (176) 243
Ripalti, A., see Battista, M.C. (176) 197
Rocco, L., Stingo, V. and Bellitti, M.
Cloning and characterization of a repetitive DNA detected by *Hin*dIII in the genome of *Raja montagui* (Batoidea, Chondrichthyes) (176) 185
Rombel, I.T., see McMorran, B.J. (176) 55
Ruiz, A., Bhat, S.P. and Bok, D.
Expression and synthesis of the Na,K-ATPase $\beta 2$ subunit in human retinal pigment epithelium (176) 237

Samuelsson, G., see Hiltonen, T. (176) 263
Sato, T., see Shimizu, E. (176) 45
Seow, H.-F., see Mertens, B. (176) 273
Shamsher, M. and Montano, X.
Analysis of intron 4 of the p53 gene in human cutaneous melanoma (176) 259
Shareck, F., see Ostiguy, S. (176) 265
Shibahara, K., see Hamada, T. (176) 211
Shimizu, E., Shirasawa, H., Kodama, K., Sato, T. and Simizu, B.
Expression, regulation and polymorphism of the *mxi1* genes (176) 45
Shirasawa, H., see Shimizu, E. (176) 45
Shirozu, M., see Hamada, T. (176) 211
Siedlecki, J.A., see Konopiński, R. (176) 191
Simizu, B., see Shimizu, E. (176) 45
Sive, H., see Kuo, J.S. (176) 17
Smith, R.G., see Neeper, M. (176) 203
Souque, P., see Chenal, V. (176) 97
Stingo, V., see Rocco, L. (176) 185
Supply, P., see Baulard, A. (176) 149

Tada, H., see Hamada, T. (176) 211
Taketo, A., see Oki, M. (176) 215
Tarassov, I., see Bordonné, R. (176) 111
Tashiro, K., see Hamada, T. (176) 211
Tatro, A., see Alves, G. (176) 39
Taura, S., see An, Z. (176) 93
Thomas, C.E., see Biegel Carson, S.D. (176) 125
Tsuchiya, E., Matsuzaki, G., Kurano, K., Fukuchi, T., Tsukao, A. and Miyakawa, T.
The *Saccharomyces cerevisiae SSD1* gene is involved in the tolerance to high concentration of Ca^{2+} with the participation of *HST1/NRC1/BFR1* (176) 35
Tsukao, A., see Tsuchiya, E. (176) 35

Valdivia, M.M., see Bolívar, J. (176) 257
Veale, R., see Kuo, J.S. (176) 17
Vidaud, D., see Baulard, A. (176) 149
Villaverde, A., see Carbonell, X. (176) 225

Walton, J.D., see Panaccione, D.G. (176) 103
Wang, S.-H., see Lee, Y.-J. (176) 85
Winkler, H., Brostjan, C., Csizmadia, V., Natarajan, G., Anrather, J. and Bach, F.H.
The intron-exon structure of the porcine E-selectin-encoding gene (176) 67

Yamada, K., see Oki, M. (176) 215

Zelenka, P.S., see Gao, C.Y. (176) 243